

MHSIP CONSUMER SURVEYS SCORING PROTOCOL

**NRI Workshop on Consumer Surveys
Arlington, Virginia
June 19-20, 2008**

MHSIP SURVEY SCORING PROTOCOL

Modifying the MHSIP Surveys

It is recommended that the surveys be used as presented (official wording, official order) for 2 reasons:

- ❑ Modifying these items may have an impact on the psychometric properties of the surveys
- ❑ Benchmarking

ADULT MHSIP SURVEY SCORING PROTOCOL

Response Scale: 1-5, Not Applicable

1 = Strongly Disagree

5 = Strongly Agree

1. Use actual scores associated with ratings
2. Recode “Not Applicable” ratings as missing
3. Exclude respondents with one-third or more missing items in the domain
4. Calculate the mean score (average) for each respondent
5. Calculate the percent of scores less than 2.5

MHSIP YOUTH SURVEY SCORING PROTOCOL

Response Scale: 1-5, Not Applicable

1 = Strongly Disagree

5 = Strongly Agree

1. Use actual scores associated with ratings
2. Recode “Not Applicable” ratings as missing
3. Exclude respondents with one-third or more missing items in the domain
4. Calculate the mean score (average) for each respondent
5. Calculate the percent of scores greater than 3.5

MHSIP SURVEY SCORING PROTOCOL

Reporting Response Rates

Indicator:
$$\frac{\text{Percent of Positive Responses}}{\text{Total Number of Responses}}$$

Response rates should be reported for each domain including Social Connectedness and Functioning

MHSIP SURVEY SCORING PROTOCOL

Recommendation:

Partition data by demographic, clinical and other variables where possible

MHSIP SURVEY SCORING PROTOCOL

Reporting National Outcome Measures

Uniform Reporting System (URS) Tables

Tables 9C, 11, 19A, 19B